arXiv:1406.0111v2 [physics.data-an] 8 May 2016

Rényi entropy and complexity measure for skew-gaussian distributions and related families

Javier E. Contreras-Reyes[a,b,*]

[a] *Division of Fisheries Research, Fisheries Development Institute, Blanco 839, Valparaíso, Chile*
[b] *Department of Mathematics, Universidad Técnica Federico Santa María, Valparaíso, Chile*

Abstract

In this paper, we provide the Rényi entropy and complexity measure for a novel, flexible class of skew-gaussian distributions and their related families, as a characteristic form of the skew-gaussian Shannon entropy. We give closed expressions considering a more general class of closed skew-gaussian distributions and the weighted moments estimation method. In addition, closed expressions of Rényi entropy are presented for extended skew-gaussian and truncated skew-gaussian distributions. Finally, additional inequalities for skew-gaussian and extended skew-gaussian Rényi and Shannon entropies are reported.

Keywords: skew-gaussian; Rényi entropy; complexity; weighted moments; Jensen's inequality

1. Introduction

The family of skew-gaussian distributions has been popularized by Azzalini (1985) and ever since it has been discussed extensively in the literature. Such discussions include a wide variety of skewed models in addition to having gaussian distribution as a special case and flexibility in capturing skewness in the data (Azzalini and Dalla-Valle 1996; Azzalini and Capitanio 1999; Azzalini 2013). In this sense, González-Farías et al. (2004) present the closed skew-gaussian distribution as an extension of the skew-gaussian case, but closed under operations such as sums, marginalization, and linear conditioning (Rezaie et al. 2014). Another generalization of the skew-gaussian distribution is the extended skew-gaussian distribution (Capitanio et al. 2003) that adds a fourth real parameter to accommodate both skewness and heavy tails. In some cases where observed variables can be simultaneously skewed and restricted to a fixed interval, the truncated skew-gaussian distribution is a good choice for those applications, especially for environmental and biological variables in which the observations are positives (Flecher et al. 2010).

In many applications, the empirical distribution of some observed variables was modeled by a skew-gaussian distribution. For example, the closed skew-gaussian distribution is used by Rezaie et al. (2014)

*Corresponding author, Phone +56 032 2151 513
Email address: jecontrr@uc.cl, javier.contreras@ifop.cl (Javier E. Contreras-Reyes)

to simulate seismic amplitude variations. Contreras-Reyes and Arellano-Valle (2012) consider the skew-gaussian distribution for seismic magnitudes of aftershocks catalogue of the 2010 Maule earthquake in Chile; Arellano-Valle et al. (2013) for the optimization of ozone's monitoring network; and Figiel (2014) for a digital reconstruction of nanocomposite morphologies from TEM (Transmission Electron Microscopy) images. An implementation of the extended skew-gaussian (in logarithmic form) can be found in Zhou and Wang (2008) for pricing of both Asian and basket options. As mentioned above, Flecher et al. (2010) considers the truncated skew-gaussian distribution to fit the daily relative humidity measurements. See more applications in Genton (2004).

More recently, Contreras-Reyes and Arellano-Valle (2012) and Arellano-Valle et al. (2013) compute the Kullback-Leibler divergence measure for skew-gaussian distribution and Shannon entropy for the full class of skew-elliptical distributions, respectively. They highlight that the Kullback-Leibler information measure should be represented in quadratic form, including a non-analytical expected value. In addition, they gave the Kullback-Leibler divergence of a multivariate skew-gaussian distribution with respect to multivariate gaussian distribution. Information measure applications dealing with skewed data have been performed by Contreras-Reyes and Arellano-Valle (2012), Arellano-Valle et al. (2013), Contreras-Reyes (2014) and references therein.

In this work, we focus on the Rényi entropy (Rényi 1970) as a characteristic form of the Shannon entropy to give a closed expression of skew-gaussian densities. Additionally, the LMC complexity measure (López-Ruiz et al. 1995) is derived by the difference between the extensive Rényi entropy and Shannon entropy (Yamano 2004). To do this, we briefly describe the main properties of closed skew-gaussian distributions. Finally, we compute the Rényi entropy and complexity measure for the extended skew-gaussian and univariate truncated skew-gaussian densities.

2. Rényi entropy and complexity measure

Consider the αth-order Rényi entropy (Rényi 1970) of probability density $f(\mathbf{x})$ on a variable $\mathbf{x} \in \Delta \subset \mathbb{R}^d$:

$$R_\alpha[f] = \frac{1}{1-\alpha} \ln \int [f(\mathbf{x})]^\alpha d\mathbf{x}, \tag{1}$$

where normalization to unity as given by $\int f(\mathbf{x}) d\mathbf{x} = 1$ (Sánchez-Moreno et al. 2014). Golshani and Pasha (2010) provide some important properties of the Rényi entropy: 1. $R_\alpha[f]$ can be negative, 2. $R_\alpha[f]$ is invariant under a location transformation, 3. $R_\alpha[f]$ is not invariant under a scale transformation, and 4. for any $\alpha_1 < \alpha_2$, $\mathbf{x} \in \Delta$, we have $R_{\alpha_1}[f] \geq R_{\alpha_2}[f]$, which are equal if and only if $\mathbf{x}$ is uniformly distributed.

From (1), the Shannon entropy is obtained by the limit

$$S[f] = \lim_{\alpha \to 1} R_\alpha[f] = -\int f(\mathbf{x}) \ln f(\mathbf{x}) d\mathbf{x} \tag{2}$$

by applying l'Hôpital's rule to $R_\alpha[f]$ with respect to α (Rényi 1970). This measure is the expected value of $g(\mathbf{x}) = -\ln f(\mathbf{x})$ with respect to $f(\mathbf{x})$, i.e., $S[f] = \langle g(\mathbf{x}) \rangle$ (Liu et al. 2012). Hereafter, we will refer to this as the expected information of $g(\mathbf{x})$ in $\mathbf{x}$. See Cover and Thomas (2006) for additional properties of the Shannon entropy.

Example 1. *(Dembo et al. 1991; Cover and Thomas 2006). Let* $\mathbf{x}$ *be gaussian with mean vector* $\mu \in \mathbb{R}^d$ *and* $\mathbf{J}$ *is a* $d \times d$ *variance matrix (with determinant* $|\mathbf{J}| > 0$*). Then, the Rényi and Shannon entropies of* $\mathbf{x}$ *are given by*

$$R_\alpha[f] = \frac{1}{2}\ln[(2\pi)^d|\mathbf{J}|] + \frac{d\ln\alpha}{2(\alpha-1)}, \quad 1 < \alpha < \infty, \tag{3}$$

$$S[f] = \frac{1}{2}\ln[(2\pi e)^d|\mathbf{J}|], \tag{4}$$

respectively.

Another important concept is the statistical complexity that measures the randomness and structural correlations of a known system (Carpi et al. 2011). López-Ruiz et al. (1995) proposed a measure of statistical complexity (LMC) in order to determine the *disequilibrium* of the system attributed to entropy measure (Anteneodo and Plastino 1996; Sánchez-Moreno et al. 2014). LMC measure is defined as the product

$$C_{LMC}[f] = e^{S[f]-R_2[f]}, \tag{5}$$

where $R_2[f]$ is the quadratic Rényi entropy of $\mathbf{x}$ ($\alpha = 2$). Yamano (2004) provide an extensive entropy instead of an additive Shannon entropy in (5), characterised as a difference between the αth-order Rényi entropy and quadratic Rényi entropy as

$$C_\alpha[f] = e^{R_\alpha[f]-R_2[f]}. \tag{6}$$

Note that $C_\alpha[f]$ reflects the shape of the distribution of $\mathbf{x}$ and takes unity for all distributions when $\alpha = 2$. In addition, $\mathcal{C}_\alpha$ satisfies a great variety of interesting mathematical and physical properties. Let us just recall here the following properties: 1. $C_\alpha[f] > 1$, $\forall\, \alpha \leq 2$, and, $0 < C_\alpha[f] \leq 1$, $\forall\, \alpha > 2$; 2. $C_\alpha[f]$ is invariant under a location and scale transformation in the distribution of $\mathbf{x}$; and 3. is invariant under replications of the original distribution of $\mathbf{x}$.

3. Skew-gaussian distribution and related families

The closed skew-gaussian distribution has interesting properties inherited from the Gaussian distribution and corresponds to a generalization of the skew-gaussian distribution. We briefly describe some of its inferential properties and present the weighted moments method in Proposition 2 (Flecher et al. 2009), necessary

to calculate Rényi entropy of skew-gaussian random vectors.

3.1. Closed skew-gaussian distributions

Concerning the definition of Flecher et al. (2009) and González-Farías et al. (2004), let $\mathbf{y} \in \Delta \subset \mathbb{R}^d$ be a random vector with closed skew-gaussian distribution denoted as $CSN_{d,s}(\mu, \mathbf{J}, \mathbf{D}, \nu, \mathbf{A})$ and with density function

$$f_{d,s}(\mathbf{y}) = \phi_d(\mathbf{y}; \mu, \mathbf{J}) \frac{\Phi_s(\mathbf{D}^\top(\mathbf{y} - \mu); \nu, \mathbf{A})}{\Phi_s(\mathbf{0}; \nu, \mathbf{A} + \mathbf{D}^\top \mathbf{J} \mathbf{D})}, \tag{7}$$

where $\mu \in \mathbb{R}^d$, $\nu \in \mathbb{R}^s$, $\mathbf{J} \in \mathbb{R}^{d \times d}$ and $\mathbf{A} \in \mathbb{R}^{s \times s}$ are both covariance matrices, $\mathbf{D} \in \mathbb{R}^{d \times s}$, $\mathbf{D}^\top$ denotes the transposed $\mathbf{D}$ matrix,

$$\phi_d(\mathbf{y}; \mu, \mathbf{J}) = \frac{1}{(2\pi)^{d/2} |\mathbf{J}|^{1/2}} \exp\left(-\frac{1}{2} (\mathbf{y} - \mu)^\top \mathbf{J}^{-1} (\mathbf{y} - \mu) \right)$$

and $\Phi_d(\mathbf{y}; \mu, \mathbf{J})$ are the probability function (pdf) and cumulative distribution function, respectively, of the d-dimensional gaussian distribution with mean vector μ and variance matrix $\mathbf{J}$. The closed skew-gaussian distribution is closed under translations, scalar multiplications, and full, row rank linear transformations (González-Farías et al. 2004; Genton 2004). Let $\mathbf{T} \in \mathbb{R}^{n \times d}$ be a matrix with rank n such that $d \leq n$, then

$$\mathbf{T}\mathbf{y} = CSN_{n,s}(\mathbf{T}\mu, \tilde{\mathbf{J}}, \tilde{\mathbf{D}}, \nu, \tilde{\mathbf{T}}) \tag{8}$$

where $\tilde{\mathbf{J}} = \mathbf{T}^\top \mathbf{J} \mathbf{T}$, $\tilde{\mathbf{D}} = \mathbf{D}^\top \mathbf{J} \mathbf{T} \tilde{\mathbf{J}}^{-1}$, and $\tilde{\mathbf{T}} = \mathbf{A} + \mathbf{D}^\top \mathbf{J} \mathbf{D} - \tilde{\mathbf{D}}^\top \tilde{\mathbf{J}} \tilde{\mathbf{D}}$ (Genton 2004, see Proposition 2.3.1).

A particular case of (8), is the standardised random vector $\mathbf{z}_0 = \mathbf{J}^{-1}(\mathbf{y} - \mu)$. In this case, Eq. (7) is rewritten as

$$f_{d,s}(\mathbf{z}_0) = \phi_d(\mathbf{z}_0) \frac{\Phi_s(\mathbf{D}^\top \mathbf{J}^{1/2} \mathbf{z}_0; \nu, \mathbf{A})}{\Phi_s(\mathbf{0}; \nu, \mathbf{A} + \mathbf{D}^\top \mathbf{J} \mathbf{D})}. \tag{9}$$

Given that the closed skew-gaussian distribution is closed under translations and by property (8), the standardised random vector $\mathbf{Z}_0$ follows $CSN_{d,s}(\mathbf{0}, \mathbf{I}_d, \mathbf{D}^\top \mathbf{J}^{1/2}, \nu, \mathbf{A})$, where $\mathbf{I}_d$ denotes the d-dimensional identity matrix. For more details, see Flecher et al. (2009) and Genton (2004).

Lemma 1. *(Flecher et al. 2009). Let* $\mathbf{Y}$ *be a* $CSN_{d,s}(\mu, \mathbf{J}, \mathbf{D}, \mathbf{0}, \mathbf{A})$*,* r *a positive integer and* $h(\mathbf{y}) = h(y_1, \ldots, y_d)$ *be any real valued function such that* $\langle h(\mathbf{Y}) \rangle$ *is finite, then*

$$\langle h(\mathbf{Y}) [\Phi_d(\mathbf{Y}; \mathbf{0}, \mathbf{I}_d)]^r \rangle = \langle h(\tilde{\mathbf{Y}}) \rangle \frac{\Phi_{rd+s}(\mathbf{0}; \tilde{\nu}, \tilde{\mathbf{A}} + \tilde{\mathbf{D}}^\top \mathbf{J} \tilde{\mathbf{D}})}{\Phi_s(\mathbf{0}; \mathbf{0}, \mathbf{A} + \mathbf{D}^\top \mathbf{J} \mathbf{D})}, \tag{10}$$

where $\tilde{\mathbf{Y}} \sim CSN_{d,rd+s}(\mu, \mathbf{J}, \tilde{\mathbf{D}}, \tilde{\nu}, \tilde{\mathbf{A}})$ *with* $\tilde{\mathbf{D}} = (\mathbf{E}^\top, \mathbf{D}^\top)$*,* $\mathbf{E}$ *a* $d \times rd$ *matrix defined by* $\mathbf{E} = (\mathbf{I}_d, \ldots, \mathbf{I}_d)$*,*

$\tilde{\nu} = (-\mu, \ldots, -\mu, \mathbf{0}_s)$ *a* $(rd + s)$ *vector and*

$$\tilde{\mathbf{A}} = \begin{pmatrix} \mathbf{I}_{rd} & \mathbf{0} \\ \mathbf{0} & \mathbf{A} \end{pmatrix}.$$

3.2. Skew-gaussian distribution

A special case of closed skew-gaussian is the gaussian density when $\mathbf{D} = \mathbf{0}$. When $s = 1$, the skew-gaussian density function is obtained (Azzalini and Dalla-Valle 1996; Azzalini and Capitanio 1999; Azzalini 2013). For simplicity, a slight variant of the original definition is considered here. In this work it is posited that a random vector $\mathbf{Z} \in \Delta \subset \mathbb{R}^d$ has a skew-gaussian distribution with mean vector $\mu \in \mathbb{R}^d$, variance matrix $\mathbf{J} \in \mathbb{R}^{d \times d}$ and shape/skewness parameter $\boldsymbol{\eta} \in \mathbb{R}^d$, denoted by $\mathbf{Z} \sim SN_d(\mu, \mathbf{J}, \boldsymbol{\eta})$, if its probability density function is

$$f(\mathbf{z}) = 2\phi_d(\mathbf{z}; \mu, \mathbf{J})\Phi_1[\boldsymbol{\eta}^\top(\mathbf{z} - \mu)]. \tag{11}$$

The mean vector and the variance matrix of $\mathbf{Z}$ are

$$\begin{aligned} \langle \mathbf{z} \rangle &= \mu + \sqrt{\frac{2}{\pi}}\,\delta, \\ \langle \mathbf{z}^2 \rangle &= \mathbf{J} - \frac{2}{\pi}\delta\delta^\top, \end{aligned}$$

respectively, where $\delta = \mathbf{J}\boldsymbol{\eta}/\sqrt{1 + \boldsymbol{\eta}^\top \mathbf{J}\boldsymbol{\eta}}$ (Azzalini and Capitanio 1999; Contreras-Reyes and Arellano-Valle 2012).

Proposition 1. *Let* $\mathbf{Z}$ *be a* $SN_d(\mu, \mathbf{J}, \boldsymbol{\eta})$*. Then:*

$$\int [f(\mathbf{z})]^\alpha d\mathbf{z} = \psi_{\alpha,d}(\mathbf{J})\,\frac{\Phi_{\alpha+1}(\mathbf{0}; \mathbf{0}, \tilde{\mathbf{J}})}{\Phi_1(0; 0, \sigma^2)}, \quad \alpha \in \mathbb{N},\ \alpha > 1, \tag{12}$$

where

$$\psi_{\alpha,d}(\mathbf{J}) = \frac{2^\alpha}{\alpha^{d/2}}[(2\pi)^d |\mathbf{J}|]^{(1-\alpha)/2},$$

$\tilde{\mathbf{J}} = \mathbf{I}_{\alpha+1} + \|\tilde{\boldsymbol{\eta}}\|^2 \tilde{\mathbf{D}}^\top \tilde{\mathbf{D}}$, $\tilde{\mathbf{D}} = (\mathbf{1}_\alpha, \|\tilde{\boldsymbol{\eta}}\|)^\top$, $\mathbf{1}_\alpha$ *is the* α*-dimensional vector of ones,* $\sigma^2 = 1 + \|\tilde{\boldsymbol{\eta}}\|^4$, $\|\tilde{\boldsymbol{\eta}}\| = \tilde{\boldsymbol{\eta}}^\top \tilde{\boldsymbol{\eta}}$ *and* $\tilde{\boldsymbol{\eta}} = \alpha^{-1/2}\mathbf{J}^{1/2}\boldsymbol{\eta}$.

By (1) and (12), the Rényi entropy of a random variable $\mathbf{Z} \sim SN_d(\mu, \mathbf{J}, \boldsymbol{\eta})$ is retrieved. Taking $\boldsymbol{\eta} = \mathbf{0}$ in (12), the Rényi entropy of the gaussian distribution given by (3) is obtained. Lemma 1 allows the computing of the expected value of the cumulative density function of a gaussian density. Considering the standarised closed skew-gaussian variable in (9), the Proposition 1 is solved by (10), by setting $\nu = \mathbf{0}$ and $\mathbf{A} = \mathbf{I}_d$, with $d = s = 1$. However, the case $\nu \neq \mathbf{0}$ and $\mathbf{A} \neq \mathbf{I}_d$, $d > 1$, is still an open problem and, it is useful to find

the Rényi entropy for closed skew-gaussian distributions. By (1) and (9), the Shannon entropy for closed skew-gaussian distributions is rewritten as

$$\begin{aligned} S[f] &= -\langle \ln[f_{d,s}(\mathbf{Y})]\rangle \\ &= \frac{1}{2}\ln|\mathbf{J}| - \ln[\Phi_s(\mathbf{0};\nu,\mathbf{A}+\mathbf{D}^\top\mathbf{J}\mathbf{D})] - \langle \ln[\phi_d(\mathbf{Z}_0)\Phi_s(\tilde{\mathbf{D}}^\top\mathbf{Z}_0;\nu,\mathbf{A})]\rangle \\ &= S[f_0] - \ln[\Phi_s(\mathbf{0};\nu,\mathbf{A}+\mathbf{D}^\top\mathbf{J}\mathbf{D})] - \langle \ln[\Phi_s(\tilde{\mathbf{D}}^\top\mathbf{Z}_0;\nu,\mathbf{A})]\rangle, \end{aligned} \tag{13}$$

where f_0 is the standardised gaussian distribution and $S[f_0] = (1/2)\ln(2\pi e)$.

Corollary 1. *Let* $\mathbf{Z} \sim SN_d(\mu,\mathbf{J},\boldsymbol{\eta})$, $\mathbf{Z}_N \sim N_d(\mu,\mathbf{J})$, $\|\tilde{\boldsymbol{\eta}}\| = \tilde{\boldsymbol{\eta}}^\top\tilde{\boldsymbol{\eta}}$ *and* $\tilde{\boldsymbol{\eta}} = \mathbf{J}^{1/2}\boldsymbol{\eta}$. *Then,*

(i) $R_\alpha[f] = R_\alpha[f_0] - N_\alpha[f], \quad \alpha \in \mathbb{N}$, $\alpha > 1$, *where*

$$N_\alpha[f] = \frac{1}{\alpha-1}\ln\left[2^\alpha \frac{\Phi_{\alpha+1}(\mathbf{0};\mathbf{0},\tilde{\mathbf{J}})}{\Phi_1(0;0,\sigma^2)}\right]$$

is the so-called Negentropy, $R_\alpha[f_0]$ *is given by (3), and* $\tilde{\mathbf{J}}$ *and* σ^2 *are defined as in Proposition 1.*

(ii) $\lim_{\alpha\to 1} N_\alpha[f] = \langle \ln[2\Phi_1(\|\tilde{\boldsymbol{\eta}}\|W)]\rangle$.

(iii) $S[f] = S[f_0] - \langle \ln[2\Phi_1(\|\tilde{\boldsymbol{\eta}}\|W)]\rangle$,

where $S[f_0]$ *is given by (4) and* $W \sim SN_1(0,1,\|\tilde{\boldsymbol{\eta}}\|)$.

(iv) $S[f_0] - \ln(4e) \le S[f] \le S[f_0]$, $\forall\,\boldsymbol{\eta}$.

Contreras-Reyes and Arellano-Valle (2012) define the negentropy as the departure from gaussianity of the distribution of $\mathbf{Z}$. Therefore, the skew-gaussian Rényi entropy corresponds to the difference between gaussian Rényi entropy and negentropy, that depends on the skewness parameter $\boldsymbol{\eta}$. On the another hand, by setting $\nu = \mathbf{0}$ and $\mathbf{A} = \mathbf{I}_d$ in (13) with $d = s = 1$, we obtain the property (ii) of Corollary 1.

By properties (iii) and (iv),

$$-0.967 \le S[f_0] - \log(4e) \le S[f]$$

because, the minimum value of normal Shannon entropy is obtained for $d = 1$ and,

$$0 \le \langle \ln[2\Phi_1(\|\tilde{\boldsymbol{\eta}}\|W)]\rangle \le 2.386,$$

for all $\boldsymbol{\eta}$. In addition, Contreras-Reyes and Arellano-Valle (2012) reported a maximum value of this expected value equal to 2.339, using numerical approximations. Considering (1), (6) and (12); the complexity measure for skew-gaussian distribution is obtained.

3.3. Extended skew-gaussian distributions

Consider a slight variant of the extended skew-gaussian distribution proposed by Capitanio et al. (2003). Let $\mathbf{Z} \sim ESN_d(\mu, \mathbf{J}, \boldsymbol{\eta}, \tau)$, $\mathbf{Z} \in \Delta \subset \mathbb{R}^d$, with mean vector $\mu \in \mathbb{R}^d$, variance matrix $\mathbf{J} \in \mathbb{R}^{d\times d}$, shape/skewness parameter $\boldsymbol{\eta} \in \mathbb{R}^d$, extended parameter $\tau \in \mathbb{R}$, and with pdf given by:

$$p(\mathbf{z}) = \frac{1}{\Phi_1(\tau)}\phi_d(\mathbf{z};\mu,\mathbf{J})\Phi_1[\boldsymbol{\eta}^\top(\mathbf{z}-\mu)+\tilde{\tau}], \tag{14}$$

where $\mathbf{z} \in \mathbb{R}^d$ and $\tilde{\tau} = \tau\sqrt{1+\boldsymbol{\eta}^\top\mathbf{J}\boldsymbol{\eta}}$. The mean vector and the variance matrix of $\mathbf{Z}$ are

$$\langle \mathbf{z} \rangle = \mu + \delta\zeta_1(\tau), \tag{15}$$

$$\langle \mathbf{z}^2 \rangle = \mathbf{J} - \zeta_1(\tau)[\tau+\zeta_1(\tau)]\delta\delta^\top, \tag{16}$$

respectively; where $\zeta_1(\mathbf{z}) = \phi(\mathbf{z})/\Phi_1(\mathbf{z})$ is the *zeta* function (Azzalini and Capitanio 1999; Capitanio et al. 2003).

Proposition 2. *Let* $\mathbf{Z}$ *be a* $ESN_d(\mu, \mathbf{J}, \boldsymbol{\eta}, \tau)$, $\mathbf{z} \in \mathbb{R}^d$. *Then:*

$$\int [f(\mathbf{z})]^\alpha d\mathbf{z} = \psi_{\alpha,d}(\mathbf{J})\langle\left[\frac{\Phi_1(W)}{2\Phi_1(\tau)}\right]^\alpha\rangle, \quad \alpha \in \mathbb{N},\ \alpha > 1, \tag{17}$$

where $\psi_{\alpha,d}(\mathbf{J})$ *is defined as in Proposition 1 and* $W = \tilde{\boldsymbol{\eta}}^\top\mathbf{Z}_0 + \tilde{\tau} \sim ESN_1(\tilde{\tau}, \|\tilde{\boldsymbol{\eta}}\|^2, \|\tilde{\boldsymbol{\eta}}\|, \tau)$, $\|\tilde{\boldsymbol{\eta}}\| = \tilde{\boldsymbol{\eta}}^\top\tilde{\boldsymbol{\eta}}$, *and* $\tilde{\boldsymbol{\eta}} = \mathbf{J}^{1/2}\boldsymbol{\eta}$.

Corollary 2. *Let* $\mathbf{Z} \sim ESN_d(\mu, \mathbf{J}, \boldsymbol{\eta}, \tau)$, $\mathbf{Z}_N \sim N_d(\mu, \mathbf{J})$ *and* W *are defined as in Proposition 2. Then,*

(i) $R_\alpha[f] = R_\alpha[f_0] - N_\alpha[f], \quad \alpha \in \mathbb{N},\ \alpha > 1,$

where

$$N_\alpha[f] = \frac{1}{\alpha-1}\ln\langle\left[\frac{\Phi_1(W)}{\Phi_1(\tau)}\right]^\alpha\rangle,$$

and $R_\alpha[f_0]$ *is given by (3).*

(ii) $R_\alpha[f] \le R_\alpha[f_0] + \dfrac{\alpha}{1-\alpha}\ln\left[\dfrac{\Phi_1(\tilde{\tau}+\tilde{\delta}\zeta_1(\tau))}{\Phi_1(\tau)}\right],$

where $\tilde{\delta} = \|\tilde{\boldsymbol{\eta}}\|^3/\sqrt{1+\|\tilde{\boldsymbol{\eta}}\|^4}$.

(iii) $S[f] = S[f_0] - \langle\ln\left[\dfrac{\Phi_1(W)}{\Phi_1(\tau)}\right]\rangle.$

(iv) $S[f_0] + \ln[\Phi_1(\tau)] - \Phi_1\left(\dfrac{\tilde{\tau}}{\sqrt{1+\boldsymbol{\eta}\boldsymbol{\eta}^\top}}\right) \le S[f] \le \dfrac{1}{2}\ln\left[(2\pi e)^d\left|\mathbf{J} - \zeta_1(\tau)[\tau+\zeta_1(\tau)]\delta\delta^\top\right|\right],\ \forall\,\boldsymbol{\eta}.$

(v) $\lim\limits_{\alpha\to 1} N_\alpha[f] = \langle\ln[\dfrac{\Phi_1(W)}{\Phi_1(\tau)}]\rangle.$

Pourahmadi (2007) illustrated the behaviour of $\zeta_1(\tau)$, $\tau \in \mathbb{R}$. This function is strictly decreasing for any $\tau \in \mathbb{R}$, tends to 0 when $\tau \to +\infty$, and diverge when $\tau \to -\infty$. For $\tau = 0$, the property (iv) of Corollary 2 becomes property (iii) of Corollary 1. By properties (iii) of Corollary 2 and (ii) of Corollary 1, the negentropy of an extended skew-gaussian random vector is always larger than the negentropy of a skew-gaussian random vector. Therefore, we obtain the following relationship among the Shannon entropies of gaussian ($f_0(\mathbf{z})$), skew-gaussian ($g(\mathbf{y})$), and extended skew-gaussian ($f(\mathbf{x})$) distributions: $S[f_0] \geq S[g] \geq S[f]$. Considering (1), (6) and (17); the complexity measure for extended skew-gaussian distribution is obtained.

3.4. Truncated skew-gaussian distributions

The truncated skew-gaussian pdf given by Flecher et al. (2010), consider the random variable $Z \sim SN_1(\mu, \omega, \lambda)$, $\mathbf{Z} \in \Delta \subset \mathbb{R}$, and the definition given in (11) for the case $d = 1$. Flecher et al. (2010) gives the expressions of the higher order and weighted moments of truncated skew-gaussian distributions. We also consider the following definition based on (11) for a truncated skew-gaussian random variable $W \in [a, b] \subset \mathbb{R}$, denoted by $W \sim TSN(\mu, J, \lambda)$, and with density

$$g(w) = \frac{f(w)}{[F(w)]_a^b}, \quad a < w \leq b, \tag{18}$$

where $f(z)$ is defined in (11) for $d = 1$ with $\mathbf{J} = J$, $\boldsymbol{\eta} = \lambda$; and $F(z)$ is the cumulative density function of Z with

$$[F(w)]_a^b = F(b) - F(a) = \int_a^b f(u)du.$$

The following Remark allows the computation of $[F(w)]_a^b$ in terms of the gaussian cumulative density function and a bivariate integral term.

Remark 1. *Let $Z \sim SN_1(\mu, J, \lambda)$, Owen (1956) and Azzalini (1985) gives the expressions to compute $F(z)$ as follows*

$$F(z) = 2\int_z^{-\infty}\int_{-\infty}^{\lambda s} \phi(s)\phi(t)\, dt\, ds = \Phi_1(z) - 2\int_z^{\infty}\int_0^{\lambda s} \phi(s)\phi(t)\, dt\, ds. \tag{19}$$

Then, by replacing (19) in $[F(w)]_a^b$ we obtain

$$[F(w)]_a^b = \Phi_1(b) - \Phi_1(a) - 2\int_a^b \int_0^{\lambda s} \phi(s)\phi(t)dtds.$$

Proposition 3. *Let Z, W be a $SN_1(\mu, J, \lambda)$ and $TSN_1(\mu, J, \lambda)$, respectively, $\lambda \neq 0$. Then:*

$$\int_a^b [g(w)]^\alpha dw = 2\psi_{\alpha,1}(J)\, \Phi_{\alpha+1}(\mathbf{0}; \mathbf{0}, \tilde{\mathbf{J}}) \frac{[H(v)]_{a_0}^{b_0}}{([F(z)]_a^b)^\alpha}, \tag{20}$$

where $\psi_{\alpha,1}(J)$ *is defined as in Proposition 1 with* $d = 1$ *and* $\mathbf{J} = J$*;* $\tilde{\mathbf{J}} = \mathbf{I}_{\alpha+1} + \tilde{\lambda}^2\tilde{\mathbf{D}}^\top\tilde{\mathbf{D}}$*,* $\tilde{\lambda}^2 = \omega\lambda^2/\alpha$*,* $\tilde{\mathbf{D}} = (\mathbf{1}_\alpha, \tilde{\lambda})^\top$ *and* $V \sim CSN_{1,2}(0, \tilde{\lambda}^2, \tilde{\mathbf{B}}, \mathbf{0}, \mathbf{I}_2)$ *with cumulative density function* $H(v)$*,* $\tilde{\mathbf{B}} = (1, \tilde{\lambda})^\top$*,* $a_0 = \lambda(a-\mu)/\omega$ *and* $b_0 = \lambda(b-\mu)/\omega$*.*

Remark 2. *By Lemma 2.2.1 of Genton (2004),* $H(v)$ *is easily computable by a tri-variate gaussian cumulative density function as*

$$H(v) = \frac{\Phi_3\left[\begin{pmatrix} v \\ \mathbf{0} \end{pmatrix}; \begin{pmatrix} 0 \\ \mathbf{0} \end{pmatrix}, \begin{pmatrix} \tilde{\lambda}^2 & -\tilde{\lambda}^2\tilde{\mathbf{B}} \\ -\tilde{\lambda}^2\tilde{\mathbf{B}}^\top & \mathbf{I}_2 + \tilde{\lambda}^2\tilde{\mathbf{B}}^\top\tilde{\mathbf{B}} \end{pmatrix}\right]}{\Phi_2(\mathbf{0}; \mathbf{0}, \mathbf{I}_2 + \tilde{\lambda}^2\tilde{\mathbf{B}}^\top\tilde{\mathbf{B}})},$$

where $\tilde{\lambda}$ *and* $\tilde{\mathbf{B}}$ *are defined as in Proposition 3.*

Considering (1), (6) and (20); the complexity measure for extended skew-gaussian distribution is obtained.

4. Conclusions

In this paper, we have presented some solutions to compute the Rényi entropy with discrete α-order and for a wide range of asymmetric distributions. Specifically, we find a closed expression for skew-gaussian, extended skew-gaussian, and truncated skew-gaussian distributions. Finally, additional inequalities for skew-gaussian and extended skew-gaussian entropies were reported.

Appendix

Proof of Proposition 1.

To compute the integral $\int [f(\mathbf{z})]^\alpha d\mathbf{z}$, we use the change of variables $\mathbf{J}_\alpha = \alpha^{-1}\mathbf{J}$ and $\mathbf{Z}_0 = \mathbf{J}_\alpha^{-1/2}(\mathbf{Z} - \mu)$, $\mathbf{Z}_0 \sim SN_d(\mathbf{0}, \mathbf{I}_d, \tilde{\boldsymbol{\eta}})$, $\tilde{\boldsymbol{\eta}} = \mathbf{J}_\alpha^{1/2}\boldsymbol{\eta}$. We shall use the fact that $|\mathbf{J}_\alpha| = \alpha^{-d}|\mathbf{J}|$ for d-dimensional matrices (Nielsen and Nock 2012). Then, according to Lemma 2 of Arellano-Valle et al. (2013), the integral $\int [f(\mathbf{z})]^\alpha d\mathbf{z}$ should be rewritten in terms of an expected value with respect to a standardized gaussian density as

$$\begin{aligned} \int [f(\mathbf{z})]^\alpha d\mathbf{z} &= \frac{2^\alpha}{|\mathbf{J}|^{\alpha/2}} |\mathbf{J}_\alpha|^{1/2} (2\pi)^{(1-\alpha)d/2} \langle [\Phi_1(\tilde{\boldsymbol{\eta}}^\top \mathbf{Z}_0)]^\alpha \rangle \\ &= \frac{2^\alpha}{\alpha^{d/2}} (2\pi)^{(1-\alpha)d/2} |\mathbf{J}|^{(1-\alpha)/2} \langle [\Phi_1(W)]^\alpha \rangle. \end{aligned}$$

where $W \sim SN_1(\mathbf{0}, \|\tilde{\boldsymbol{\eta}}\|^2, \|\tilde{\boldsymbol{\eta}}\|)$ with $\|\tilde{\boldsymbol{\eta}}\| = \tilde{\boldsymbol{\eta}}^\top\tilde{\boldsymbol{\eta}}$ (Contreras-Reyes and Arellano-Valle 2012; Arellano-Valle et al. 2013), i.e., the expected value $\langle [\Phi_1(\tilde{\boldsymbol{\eta}}^\top \mathbf{Z}_0)]^\alpha \rangle$ is reduced from d dimensions to one dimension (Arellano-Valle et al.

2013; Contreras-Reyes 2014). By Lemma 1 and setting $\mu = \mathbf{0}$, $\mathbf{J} = \|\tilde{\boldsymbol{\eta}}\|^2$, $\mathbf{D} = \|\tilde{\boldsymbol{\eta}}\|$, $r = \alpha$, $\mathbf{A} = s = h(w) = 1$; we obtain $\tilde{\mathbf{A}} = \mathbf{I}_{\alpha+1}$ and $\tilde{\mathbf{D}} = (\mathbf{1}_\alpha, \|\tilde{\boldsymbol{\eta}}\|)^\top$. Therefore, the expected value of the integral is reduced to

$$\langle [\Phi_1(W)]^\alpha \rangle = \frac{\Phi_{\alpha+1}(\mathbf{0}; \mathbf{0}, \mathbf{I}_{\alpha+1} + \|\tilde{\boldsymbol{\eta}}\|^2 \tilde{\mathbf{D}}^\top \tilde{\mathbf{D}})}{\Phi_1(0; 0, 1 + \|\tilde{\boldsymbol{\eta}}\|^4)}. \ \square$$

Proof of Corollary 1

(i) Follows from (3) and Proposition 1.

(ii) See Proposition 2 of Arellano-Valle et al. (2013).

(iii) Right side: see Contreras-Reyes and Arellano-Valle (2012). Left side: consider the nonsymmetrical entropy of Liu (2009) given by

$$S(\mathbf{u}) = -\int f(\mathbf{u}) \ln[\beta(\mathbf{u}) f(\mathbf{u})] \, d\mathbf{u},$$

where $f(\mathbf{u})$ is the probability density function of a gaussian variable $\mathbf{u}$. By choosing $\beta(\mathbf{u}) = 2\Phi_1(\boldsymbol{\eta}^\top \mathbf{J}^{-1/2}(\mathbf{u} - \mu))$, $\mathbf{u} = \mathbf{Z}_N$, it follows that $\langle \ln\beta(\mathbf{Z}_N) \rangle = \ln[2] + \Phi_1(0) = (1/2)\ln(4e)$ (see Proposition 4 of Azzalini and Dalla-Valle 1996). Then, as $\langle \ln\beta(\mathbf{Z}) \rangle \leq 2\langle \ln\beta(\mathbf{Z}_N) \rangle$, the result is obtained.

(iv) Follows from properties (i), (ii) and (1). $\square$

Proof of Proposition 2

By (14), $\phi_d(\mathbf{y}; \mu, \mathbf{J}) = |\mathbf{J}|^{-1/2} \phi_d\left(\mathbf{J}^{-1/2}(\mathbf{y} - \mu)\right)$, where $\phi_d(\mathbf{z})$ is the probability density function of $N_d(\mathbf{0}, \mathbf{I}_d)$. Then, as in (1), to compute the integral $\int [f(\mathbf{z})]^\alpha d\mathbf{z}$ we use the change of variables $\mathbf{J}_\alpha = \alpha^{-1}\mathbf{J}$ and $\mathbf{Z}_0 = \mathbf{J}_\alpha^{-1/2}(\mathbf{Z} - \mu)$. In this case, $\mathbf{Z}_0 \sim ESN_d(\mathbf{0}, \mathbf{I}_d, \tilde{\boldsymbol{\eta}}, \tau)$ with $\tilde{\boldsymbol{\eta}} = \mathbf{J}_\alpha^{1/2}\boldsymbol{\eta}$. We shall use the fact that $|\mathbf{J}_\alpha| = \alpha^{-d}|\mathbf{J}|$ for d-dimensional matrices (Nielsen and Nock 2012). Then, according to Lemma 2 of Arellano-Valle et al. (2013), the integral $\int [f(\mathbf{z})]^\alpha d\mathbf{z}$ should be rewritten in terms of an expected value with respect to a standardized gaussian density as

$$\begin{aligned} \int [f(\mathbf{z})]^\alpha d\mathbf{z} &= \frac{1}{[\Phi_1(\tau)]^\alpha} |\mathbf{J}|^{-\frac{\alpha}{2}} |\mathbf{J}_\alpha|^{1/2} (2\pi)^{(1-\alpha)\frac{d}{2}} \langle [\Phi_1(\tilde{\boldsymbol{\eta}}^\top \mathbf{z}_0 + \tilde{\tau})]^\alpha \rangle \\ &= \frac{1}{[\Phi_1(\tau)]^\alpha} \alpha^{-d} (2\pi)^{(1-\alpha)d/2} |\mathbf{J}|^{(1-\alpha)/2} \langle [\Phi_1(W)]^\alpha \rangle. \end{aligned}$$

where $W = \tilde{\boldsymbol{\eta}}^\top \mathbf{Z}_0 + \tilde{\tau} \sim ESN_1(\tilde{\tau}, \|\tilde{\boldsymbol{\eta}}\|^2, \|\tilde{\boldsymbol{\eta}}\|, \tau)$ with $\|\tilde{\boldsymbol{\eta}}\| = \tilde{\boldsymbol{\eta}}^\top \tilde{\boldsymbol{\eta}}$ (Contreras-Reyes and Arellano-Valle 2012; Arellano-Valle et al. 2013), i.e., the expected value $\langle [\Phi_1(\tilde{\boldsymbol{\eta}}^\top \mathbf{z}_0 + \tilde{\tau})]^\alpha \rangle$ is reduced from d dimensions to one dimension (Arellano-Valle et al. 2013; Contreras-Reyes 2014). $\square$

Proof of Corollary 2

(i) From Proposition 2, we obtain directly

$$\begin{aligned} R_\alpha[f] &= \frac{1}{1-\alpha}\left(\ln[\psi_{\alpha,d}(\mathbf{J})] - \alpha\ln[2\Phi_1(\tau)] + \ln[\langle[\Phi_1(W)]^\alpha\rangle]\right), \\ &= R_\alpha[f_0] + \frac{\alpha}{1-\alpha}\ln\left[\frac{1}{\Phi_1(\tau)}\right] + \frac{1}{1-\alpha}\ln[\langle[\Phi_1(W)]^\alpha\rangle]. \end{aligned}$$

(ii) Considering Jensen's inequality, we obtain $\langle[\Phi_1(W)]^\alpha\rangle \geq [\Phi_1(\langle W\rangle)]^\alpha$. Then, (ii) is straightforward from (15).

(iii) By (1), it follows that

$$S[f] = -\langle\ln\left[\phi_d(\mathbf{Z}_0)\frac{\Phi_1(\tilde{\boldsymbol{\eta}}^\top\mathbf{Z}_0+\tilde{\tau})}{\Phi_1(\tau)}\right]\rangle = S[f_0] - \langle\ln\left[\frac{\Phi_1(W)}{\Phi_1(\tau)}\right]\rangle,$$

where, as in Proposition 2, $\mathbf{Z}_0 = \mathbf{J}^{-1/2}(\mathbf{Z}-\mu) \sim ESN_d(\mathbf{0}, \mathbf{I}_d, \tilde{\boldsymbol{\eta}}, \tau)$ and $W = \tilde{\boldsymbol{\eta}}^\top\mathbf{Z}_0 + \tilde{\tau} \sim ESN_1(\tilde{\tau}, \|\tilde{\boldsymbol{\eta}}\|^2, \|\tilde{\boldsymbol{\eta}}\|, \tau)$.

(iv) Right side: by Cover and Thomas (2006), for any density $g(\mathbf{x})$ of a random vector $\mathbf{x} \in \Delta \subset \mathbb{R}^d$ (not necessary gaussian) with zero mean and variance $\mathbf{J} = \langle\mathbf{X}\mathbf{X}^\top\rangle$, the Shannon entropy of $\mathbf{x}$ is maximized under gaussianity as $S[g] \leq (1/2)\ln[(2\pi e)^d|\mathbf{J}|]$. Then, the result is obtained from (16). Left side: as in Corollary 1 (iii), by choosing $\beta(\mathbf{u}) = \Phi_1(\boldsymbol{\eta}^\top\mathbf{J}^{-1/2}(\mathbf{u}-\mu)+\tilde{\tau})/\Phi_1(\tau)$ in the nonsymmetrical entropy, it follows that

$$\langle\ln\beta(\mathbf{Z}_N)\rangle = \Phi_1\left(\frac{\tilde{\tau}}{\sqrt{1+\|\boldsymbol{\eta}\|}}\right) - \ln[\Phi_1(\tau)]$$

(see Proposition 4 of Azzalini and Dalla-Valle 1996). Then, as $\langle\ln\beta(\mathbf{Z})\rangle \leq \langle\ln\beta(\mathbf{Z}_N)\rangle/\Phi_1(\tau)$, the result is obtained.

(v) Follows from properties (i), (iii) and (1). □

Proof of Proposition 3

By (18), it follows that

$$\int_a^b [g(w)]^\alpha dw = \frac{1}{([F(z)]_a^b)^\alpha}\int_a^b [f(w)]^\alpha dw$$

and, by Proposition 1, the integral $\int_a^b[f(w)]^\alpha dw$ should be rewritten in terms of an expected value as

$$\int_a^b [f(w)]^\alpha dw = \psi_{\alpha,1}(J)\langle[\Phi_1(u)]^\alpha | a_0 < u \leq b_0\rangle,$$

where $U \sim SN_1(0, \tilde{\lambda}^2, \tilde{\lambda})$, $\tilde{\lambda}^2 = \omega\lambda^2/\alpha$, $a_0 = \lambda(a-\mu)/\omega$ and $b_0 = \lambda(b-\mu)/\omega$. Again, by Lemma 1 and setting $\mu = 0$, $J = \tilde{\lambda}^2$, $r = \alpha$, $d = s = \mathbf{A} = h(u) = 1$; we obtain $\tilde{\mathbf{A}} = \mathbf{I}_{\alpha+1}$, $\tilde{\mathbf{D}} = (\mathbf{1}_\alpha, \tilde{\lambda})^\top$ and $\tilde{\mathbf{J}} = \mathbf{I}_{\alpha+1} + \tilde{\lambda}^2\tilde{\mathbf{D}}^\top\tilde{\mathbf{D}}$. Then, the expected value is

$$\langle[\Phi_1(u)]^\alpha | a_0 < u \leq b_0\rangle = 2\Phi_{\alpha+1}(\mathbf{0}; \mathbf{0}, \tilde{\mathbf{J}})[H(v)]_{a_0}^{b_0},$$

where $H(v)$ is the cumulative density function of a closed skew-gaussian variable $V \sim CSN_{1,2}(0, \tilde{\lambda}^2, \tilde{\mathbf{B}}, \mathbf{0}, \mathbf{I}_2)$ with $\tilde{\mathbf{B}} = (1, \tilde{\lambda})^\top$ (see Proposition 3 of Flecher et al. 2010). □

Acknowledgements

This work was supported by Instituto de Fomento Pesquero (IFOP, `http://www.ifop.cl/`), Valparaíso, Chile. The author would like to thank the editor and an anonymous referee for their helpful comments and suggestions.

References

Azzalini, A., 1985. A Class of Distributions which includes the Normal Ones. Scand. J. Stat. 12, 171-178.

Azzalini, A., Dalla-Valle, A., 1996. The multivariate skew-normal distribution. Biometrika 83, 715-726.

Azzalini, A., Capitanio, A., 1999. Statistical applications of the multivariate skew normal distributions. J. Roy. Stat. Soc. Ser. B 61, 579-602.

Azzalini, A., 2013. The Skew-Normal and Related Families. Vol. 3, Cambridge University Press.

González-Farías, G., Domínguez-Molina, J., Gupta, A., 2004. Additive properties of skew normal random vectors. J. Stat. Plann. Inference 126, 521-534.

Rezaie, J., Eidsvik, J., Mukerji, T., 2014. Value of information analysis and Bayesian inversion for closed skew-normal distributions: Applications to seismic amplitude variation with offset data. Geophys. 79, R151-R163.

Capitanio, A., Azzalini, A., Stanghellini, E., 2003. Graphical models for skew-normal variates. Scand. J. Stat. 30, 129-144.

Flecher, C., Allard, D., Naveau, P., 2010. Truncated skew-normal distributions: moments, estimation by weighted moments and application to climatic data. Metron 68, 265-279.

Contreras-Reyes, J.E., Arellano-Valle, R.B., 2012. Kullback-Leibler divergence measure for Multivariate Skew-Normal Distributions. Entropy 14, 1606-1626.

Arellano-Valle, R.B., Contreras-Reyes, J.E., Genton, M.G., 2013. Shannon entropy and mutual information for multivariate skew-elliptical distributions. Scand. J. Stat. 40, 42-62.

Figiel, Ł., 2014. Effect of the interphase on large deformation behaviour of polymer–clay nanocomposites near the glass transition: 2D RVE computational modelling. Comput. Mater. Sci. 84, 244-254.

Zhou, J., Wang, X., 2008. Accurate closed-form approximation for pricing Asian and basket options. Appl. Stochastic Models Bus. Ind. 24, 343-358.

Genton, M.G., 2004. Skew-elliptical distributions and their applications: A journey beyond normality. Chapman & Hall/CRC, Boca Raton, FL.

Contreras-Reyes, J.E., 2014. Asymptotic form of the Kullback-Leibler divergence for multivariate asymmetric heavy-tailed distributions. Physica A 395, 200-208.

Rényi, A., 1970. Probability theory. North-Holland, Amsterdam.

López-Ruiz, R., Mancini, H.L., Calbet, X., 1995. A statistical measure of complexity. Phys. Lett. A 209, 321-326.

Anteneodo, C., Plastino, A.R., 1996. Some features of the López-Ruiz-Mancini-Calbet (LMC) statistical measure of complexity. Phys. Lett. A 223, 348-354.

Yamano, T., 2004. A statistical measure of complexity with nonextensive entropy. Physica A 340, 131-137.

Carpi, L.C., Rosso, O.A., Saco, P.M., Ravetti, M.G., 2011. Analyzing complex networks evolution through Information Theory quantifiers. Phys. Lett. A 375, 801-804.

Sánchez-Moreno, P., Angulo, J.C., Dehesa, J.S., 2014. A generalized complexity measure based on Rényi entropy. Eur. Phys. J. D 68, 212.

Golshani, L., Pasha, E., 2010. Rényi entropy rate for Gaussian processes. Inform. Sci. 180, 1486-1491.

Liu, T., Zhang, P., Dai, W-.S., Xie, M., 2012. An intermediate distribution between Gaussian and Cauchy distributions. Physica A 391, 5411-5421.

Cover, T.M., Thomas, J.A., 2006. Elements of information theory. Wiley & Son, Inc., New York, NY, USA.

Dembo, A., Cover, T.M., Thomas, J.A., 1991. Information Theoretic Inequalities. IEEE Trans. Inform. Theory 37, 1501-1518.

Flecher, C., Naveau, P., Allard, D., 2009. Estimating the Closed Skew-Normal distributions parameters using weighted moments. Stat. Prob. Lett. 79, 1977-1984.

Pourahmadi, M., 2007. Skew-Normal ARMA Models with Nonlinear Heteroscedastic Predictors. Commun. Stat. A-Theor. 36, 1803-1819.

Owen, D.B., 1956. Tables for computing bivariate normal probabilities. Ann. Math. Stat. 27, 1075-1090.

Nielsen, F., Nock, R., 2012. A closed-form expression for the Sharma–Mittal entropy of exponential families. J. Phys. A: Math. Theor. 45, 032003.

Liu, C.-S., 2009. Nonsymmetric entropy and maximum nonsymmetric entropy principle. Chaos Soliton. Fract. 40, 2469-2474.